

Modified Recording Chamber for Sustained Kainate-Induced Gamma Oscillations in Submerged Rat Hippocampal Slices

Juzekaeva E., Nasretidinov A., Mukhtarov M.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. Fast neuronal network oscillations in an in vitro model are a challenging task. Here, we report that sustained gamma (31–35 Hz) oscillations can be induced by kainate (50 nM) in submerged rat hippocampal slices using modified Hájos's type recording chamber with a superfusion inlet positioned close to the CA3 pyramidal cell layer. The general features of these kainate-induced gamma oscillations were similar to those previously reported in the hippocampal slices using the interface-type chamber and superfused hippocampus in vivo. We suggest that close positioning of the superfusion inlet improves oxygen supply and temperature control of the oscillation-generating network and that this modification could be useful in studies of the gamma rhythmogenesis in the submerged slices.

<http://dx.doi.org/10.1007/s12668-016-0313-y>

Keywords

Beta, Gamma, Hippocampus, In vitro, Kainate, Oscillations, Submerged

References

- [1] Yamamoto, C., & McIlwain, H. (1966). Potentials evoked in vitro in preparations from the mammalian brain. *Nature*, 210(5040), 1055–1056.
- [2] Steriade, M. (2001). The intact and sliced brain. Cambridge: MIT Press.
- [3] Fisahn, A., Pike, F. G., Buhl, E. H., Paulsen, O. (1998). Cholinergic induction of network oscillations at 40 Hz in the hippocampus in vitro. *Nature*, 394(6689), 186–189. doi:10.1038/28179.
- [4] Hájos, N., Katona, I., Naiem, S. S., Mackie, K., Ledent, C., Mody, I., et al. (2000). Cannabinoids inhibit hippocampal GABAergic transmission and network oscillations. *European Journal of Neuroscience*, 12(9), 3239–3249. doi:10.1046/j.1460-9568.2000.00217.x.
- [5] Khazipov, R., & Holmes, G. L. (2003). Synchronization of kainate-induced epileptic activity via GABAergic inhibition in the superfused rat hippocampus in vivo. *The Journal of Neuroscience*, 23(12), 5337–5341.
- [6] McMahon, L. L., Williams, J. H., Kauer, J. A. (1998). Functionally distinct groups of interneurons identified during rhythmic carbachol oscillations in hippocampus in vitro. *The Journal of Neuroscience*, 18(15), 5640–5651.
- [7] Hájos, N., Ellender, T. J., Zemankovics, R., Mann, E. O., Exley, R., Cragg, S. J., et al. (2009). Maintaining network activity in submerged hippocampal slices: importance of oxygen supply. *European Journal of Neuroscience*, 29(2), 319–327. doi:10.1111/j.1460-9568.2008.06577.x.
- [8] Gulyás, A. I., Szabó, G. G., Ulbert, I., Holderith, N., Monyer, H., Erdélyi, F., et al. (2010). Parvalbumin-containing fast-spiking basket cells generate the field potential oscillations induced by cholinergic receptor activation in the hippocampus. *The Journal of Neuroscience*, 30(45), 15134–15145. doi:10.1523/JNEUROSCI.4104-10.2010.

- [9] Tsintsadze, V., Minlebaev, M., Suchkov, D., Cunningham, M. O., Khazipov, R. (2015). Ontogeny of kainate-induced gamma oscillations in the rat CA3 hippocampus in vitro. *Frontiers in Cellular Neuroscience*, 9(195), 1-15. doi:10.3389/fncel.2015.00195.